

HEE CONTRIBED SHAYES OF ANTERIOR

Monsanto Technology T. L.C.

MICLERS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE **EXAMINATION** MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY WARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC KON EVISHMENT OF VIABLE BASIC SEED OF THE VARIEUT IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE EXCLUDE OTHERS FROM SELLING THE WARTETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IN OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE RPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'I006234'

In Testimone Mercest, I have hereunto set my hand and caused the seal of the Hant Hariety Hestertion Office to be affixed at the City of Washington, D.C. this twenty-ninth day of April, in the year two thousand and eight.

Commissioner Plant Variety Protection Office Colmand 4: Selves

ST-470 (02-10-2003) designed by the Plant Variety Protection Office using

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initiated and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21, CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Parent of a hybrid sold in the U.S. - March 2004

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

U.S. Patent Application No. 10/819,099 - filed April 6, 2004 (1006234)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filling a change of address. The fee for filling a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center--East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a comptaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer. ST-470 (02-10-2003) designed by the Plant Variety Protection Office with V

EXHIBIT A (revised)

Origin and Breeding History 1006234

Corn Variety I006234 was selected for greater combining ability and yield, and resistance to ear droppage.

| Winter 1997-98 | The inbred line 01HGI4 (a proprietary DEKALB Genetics Corporation inbred) was crossed to the inbred line 4KDH1* (a proprietary DEKALB Genetics Corporation inbred) in nursery rows 3H:10-9 and 3H:13-12 at Kihei, HI. The S0 seed was subsequently grown and self-pollinated in |
|------------------|---|
| Summer 1998 | nursery row 2-19 at Kihei, HI. The S1 seed was grown and self-pollinated in nursery rows 4-4 through 5-6 at Kihei, HI. 105 ears were selected. |
| Summer 1999 | S2 ears were grown ear-to-row and self-pollinated at Stonington, IL. 3 ears were selected and bulked in nursery row 285-43. |
| Winter 1999-2000 | S3 bulk seed was grown ear-to-row and self-pollinated in nursery row PRSS2-2727 at Isabela, PR. 3 ears were selected. |
| Summer 2000 | S4 ears were grown ear-to-row and self-pollinated at Stonington, IL. 3 ears from nursery row 502-52 were selected and designated as corn variety I006234 . |
| Winter 2000-01 | S5 ears were grown ear-to-row and self-pollinated at San Juan, Jalisco, Mexico in nursery rows MXSS4:1266-1268. 4 ears were selected from row 1267 and 5 ears were selected from row 1268. |
| Summer 2001 | S6 ears were grown ear-to-row and self-pollinated at Stonington, IL. Final selection was made by selecting 5 ears each from nursery rows 525:67-68 and 525:75-76 (total 10 ears). |

^{*-} Inbred line 4KDH1 is derived from public line MO17.

Statement of Stability and Uniformity

Corn variety 1006234 was coded in 2000 and has been reproduced and judged stable for the past three generations by self-pollination. Corn variety 1006234 is uniform for all traits observed.

Statement of Variants

Corn Variety I006234 shows no variants other than that would be expected due to environment or that would occur for almost any character during the course of repeated sexual reproduction.

Statement of Distinctness

Monsanto Technology LLC believes that Corn Variety I006234 is most similar to Corn Variety 6HAK02, a proprietary DEKALB Genetics Corporation inbred:

Corn Variety I006234 differ from Corn Variety 6HAK02 at the following traits:

| Trait | 1006234 | 6HAK02 |
|--------------|------------------------------|------------------------|
| Anther Color | Yellow (2.5 Y 8/10) | Pink |
| Glume Color | (2.5 f 6/10) Green | (2.5 R 7/6) |
| | (5 GY 4/8) | Light Red (5 R 5/8) |
| Silk Color | Green-Yellow (2.5 GY 8/6) | Red (2.5 R 4/8) |

Corn Variety I006234 has yellow anther color, green glume color and Green-yellow silk color while comparative corn variety 6HAK02 has pink anther color, light red glume color and red silk color.

Description of Experimental Design

The corn varieties I006234, 6HAK02 and MO17 were grown at the Waterman, IL observation nursery in years 2002-2003. The varieties were planted in 2 row plots with 15 plants per row in each of the three years. Trait data were collected on 10 random representative plants for most traits from each 2 row plot. Data on qualitative traits are usually collected on 10 plants from each 2 row plot. For Exhibit C all data were pooled and reported as means across the years for subject variety and the standard variety with standard deviation. The varieties are randomly planted in a 4.5 acre observation nursery which is located within a larger 18 acre field. Besides the observation nursery, this field consists of a research seed increase nursery and an IP seed inventory nursery. The location of each of these individual nurseries is rotated each year to a different location within the 18 acre field. Therefore subject inbreds are not planted adjacent to comparative or standard varieties and may be located in different areas of the larger field each year, therefore being influenced by spacial differences within the field. Growing conditions within the field are not uniform as there are some slight topographical variations such as lower areas which may accumulate and retain water or higher areas which are usually drier. The field is tiled and therefore a variety maybe planted close to a tile line while a comparative variety maybe planted further away and in a low spot within the field. Temporal varieties can exist as weather conditions from year to year can vary as well as planting dates can vary from year to year based on weather conditions. Weather conditions each year can vary the maturity rate of the varieties due to either favorable or unfavorable growing conditions.

Trait variability is not observed for each variety within its own test plot-plants are usually uniform and data are collected on the "most" representative plants- variability occurs due to spacial location of the test plot for that variety from year to year and to the temporal variation of weather conditions from year to year during the 2-3 years data are collected.

Waterman Research Station Weather Data 2002-2003

| Date | Average Precip. (mm) | Ave. Monthly Temp – Max. | Ave. Monthly Temp-Min | Ave. Monthly Rel. Humid | Ave. Monthly Rel. Humid – |
|-------------|----------------------------|-----------------------------|--------------------------|----------------------------|------------------------------|
| June 2002 | | (F°) | (F°) | Max (%) | Min (%) |
| | 5.3 | 81.3 | 60.4 | 90.7 | 47.7 |
| July 2002 | 1.5 | 87.0 | 64.9 | 93.2 | 48.3 |
| August 2002 | 5.7 | 83.1 | 61.0 | 96.0 | 51.8 |
| Sept. 2002 | 1.5 | 79.4 | 52.6 | 95.0 | 42.7 |
| June 2003 | 1.7 | 76.0 | 54.0 | 90.6 | 44.3 |
| July 2003 | 3.3 | 82.0 | 60.0 | 93.6 | 53.2 |
| August 2003 | 1.3 | 84.0 | 61.0 | 93.0 | 50.5 |
| Sept 2003 | 2.1 | 74.0 | 51.0 | 92.4 | 42.9 |

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

| Name of Applicant(s) | | Variety Seed Sou | irce | Variety | Name or Ter | mporary Designation |
|--|---|---|---------------------|----------------------------|--|--|
| Monsanto Technology L.L.C. | | variety beda boo | | vullely | | 6234 |
| Address (Street & No., or R.F.D. No., City, State, Zip Code and Country) FOR OFFICIAL USE | | | | | | |
| • | ma Country) | | - | FOR OFFI | CIAL USE | |
| 800 N. Lindbergh Blvd. Creve Coeur, MO U.S.A. | | | | PVPO Numi | | 500157 |
| Place the appropriate number that describes the varietal characters typical of this inbred variety in the spaces below. Right j whole numbers by adding leading zeroes if necessary. Completeness should be striven for to establish an adequate variety descri Traits designated by a '*' are considered necessary for an adequate variety description and must be completed. | | | | | | low. Right justify ariety description. |
| 03=Dark Green 08=Yellow-Orange 13=C 04=Very Dark Green 09=Salmon 14=R | Pink Light Red Cherry Red | 16=Pale E 17=Purple 18=Colorl 19=White | Purple e less | 2 2 2 2 2 2 | 1=Buff 2=Tan 3=Brown 4=Bronze | ed (Describe) |
| STANDARD INBRED CHOICES(Use the most similar (in background a Yellow Dent Families: Family Members | of these to makent (Unrelated): 09, ND246, | ke compa | Sweet | Corn: | w-out trial data): 125, P39, 2132 | |
| B14 CM105, A632, B64, B68 B37 B37, B76, H84 B73 N192, A679, B73, NC268 C103 Mo17, Va102, Va35, A682 Ch43 A619, MS71, H99, Va26 WF9 W64A, A554, A654, Pa91 | Oh7, T2: W117, W1 W182BN White Der C166, H1 | L53R | | Pipec | 33, 4722, | HP301, HP7211 Mo24W |
| 1. TYPE: (describe intermediate types in Comments section) * 2 1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental | 7=Pipecorr | n | Stand 2 | ard Inbre | d Name MO1 | 7 |
| 2. REGION WHERE DEVELOPED IN THE U.S.A.: * 2 1=Northwest 2=Northcentral 3=Northeast 4=Southeast 6=Southwest 7=Other | 5=Southcent | ral | Stand 2 | ard Seed | Source NCR | IPS_ |
| 3. MATURITY (In Region Best Adaptability; show Heat Unit form section): DAYS HEAT UNITS * _ 8 4 | plants in si plants in po ed edible quali | .lk bllen ty | | 9 5 | HEAT UN: 1 6 8 1 5 8 | 0. 0 |
| 4. PLANT: Standar | rd Deviation | Sample Size | | Sta | ndard Devi | ation Sample Size |
| * 2 2 7.4 cm Plant Height (to tassel tip) | 8.2 | 20 | 1 9 | 2. 7 | 18.6 | 30 |
| * 0 9 2.5 cm Ear Height (to base of top ear node) | 0.7 | 20 | 0 7 | 6.8 | 14.0 | 30 |
| 0 1 4.1 cm Length of Top Ear Internode | 1.9 | 20 | 0 1 | 4.4 | 1.8 | 30 |
| _ Average Number of Tillers | | | | | | |
| * 2. 0 Average Number of Ears per Stalk | 0.7 | 20 | 0 0 | 1. 0 | 0.0 | 30 |
| 1 Anthocyanin of Brace Roots: 1=Absent 2=Faint 3=Moderate 4=Dark | | | | | | |
| Application Variety Data | Pag | e 1 | Stand | ard Inbre | d Data | |

| Application Variety Data | Page 2 Standard Inbred Data | | | , | |
|--|-----------------------------|-------------|--------------|-------------------|---------------|
| 5. LEAF: | Standard Deviation | Sample Size | s | tandard Deviation | n Sample Size |
| * 0 0 8.8 cm Width of Ear Node Leaf | 0.5 | 20 | 0 0 9.0 | 0.7 | 30 |
| * 0 7 8.6 cm Length of Ear Node Leaf | 4.2 | 20 | 0 6 2.4 | 6.4 | 30 |
| * 5. 4 Number of leaves above top ear | 0.5 | 15 | 5. 6 | 0.4 | 15 |
| 1 9. 5 degrees Leaf Angle | 3.2 | 20 | 3 5.8 | 7.8 | 30 |
| <pre>(measure from 2nd leaf above ear at * 0 3 Leaf Color (Munsell code 5 GY 3/4)</pre> | anthesis to stalk abo | ve leaf) | | | |
| 3 Leaf Sheath Pubescence(Rate on scal | o from 1 +- 0 | -l. C) | , , | ll code 5 GY 5/10 | 0) |
| 5 Marginal Waves (Rate on scale from | | cn iuzz) | 2 | | |
| 7 Longitudinal Creases (Rate on scale | | , | 5 8 | | |
| 6. TASSEL: | | | | | |
| | Standard Deviation | Sample Size | | tandard Deviation | • |
| * 0 4. 9 Number of Primary Lateral Branches 3 1. 3 Branch Angle from Central Spike | 1.3 | 20 | 7. 1 | 1.1 | 30 |
| * 4 1. 6 cm Tassel Length | 9.7 | 20 | 3 4.6 | 5.2 | 30 |
| (from top leaf collar to tassel tip) 4. 4 Pollen Shed (Rate on scale from 0≔male s | 4.5 | 20 | 4 7.4 | 4.9 | 30 |
| 0 7 Anther Color (Munsell code 2.5 Y 8/10) | cerite to 9=neavy sned | , | 4.3 | 11 1- 0 F OV O | 15) |
| 0 2 Glume Color (Munsell code 5 GY 4/8) | | | · | 11 code 2.5 GY 8, | |
| 1 Bar Glumes (Glume Bands): 1=Absent 2=Pre: | sent | | 0 2 (Munse | 11 code 5 GY 4/8; | 1 |
| | Serie | | 1 | | |
| 7a. EAR (Unhusked Data): | | | 0 5 00 | 11 2 f CV 0 | (6) |
| * 0 5 Silk Color (3 days after emergence) (Munse: | ll code 2.5 GY 8/6) | | | ll code 2.5 GY 8, | |
| 0 2 Fresh Husk Color (25 days after 50% silking | g) (Munsell code 5 GY | 1/8) | | ll code 5 GY 4/8; | |
| 2 1 Dry Husk Color (65 days after 50% Silking) | (Munsell code 2.5 Y 8, | 4) | 2 I (Munse. | ll code 2.5 Y 8/4 | 4) |
| * 3 Position of Ear at Dry Husk Stage: 1=Upright | nt 2=Horizontal 3=Pende | ent | 8 | | |
| 7 Husk Tightness (Rate on scale from 1=very) | loose to 9=very tight) | | 3 | | |
| 2 Husk Extension (at harvest): 1=Short (ears 3 =Long (8-10 cm beyond ear | | | 3 | | |
| | exp, recry being (>10 | City | | | |
| 7b. EAR (Husked Ear Data): | Standard Deviation | Sample Size | St | tandard Deviation | ı Sample Size |
| * 1 3.7 cm Ear Length | 0.9 | 20 | 1 8.5 | 0.7 | 30 |
| * 4 0.9 mm Ear Diameter at mid-point | 1.8 | 20 | 3 8.0 | 1.6 | 30 |
| 1 0` 1. 4 gm Ear Weight | 6.7 | 20 | 1 0 4.8 | 18.0 | 30 |
| * 1 3 Number of Kernel Rows | 1.1 | 15 | 1 2 | 0.7 | 15 |
| 2 Kernel Rows: 1=Indistinct 2=Distinct | | | 2 | | |
| 1 Row Alignment: 1=Straight 2=Slightly C | Curved 3=Spiral | | 1 | | |
| 0 9.3 cm Shank Length | 0.8 | 20 | 0 9.8 | 1.9 | 15 |
| 2 Ear Taper: 1=Slight 2=Average 3=Extrem | ie | | 2 | | |
| Application Variety Data | | | Standard Inb | red Data | |
| Note: Use chart on first page to choose color codes f | or color traits. | | | | |

| | | · · · · · · · · · · · · · · · · · · · | 1 | ~ ~ | _ , - , |
|--|--|---------------------------------------|--|-----------------------|-------------|
| Application Variety Data | Page 3 | | Standard Inbred Data | | |
| 8. KERNEL (Dried): | Standard Deviation | Sample Size | | Standard Deviation | Sample Size |
| 1 0.6 mm Kernel Length | 0.5 | 15 | 1 1.4 | 0.4 | 15 |
| 0 8.4 mm Kernel Width | 0.4 | 15 | 0 9.0 | 0.5 | 15 |
| 0 3.9 mm Kernel Thickness | 0.4 | 15 | 0 4.9 | 0.3 | 15 |
| 2 2.4 % Round Kernels (Shape Grade) | 1.8 | 15 | 3 1 .7 | 3.6 | 15 |
| 1 Aleurone Color Pattern: 1=Homozygous 2=Se | · · | | 1 | | |
| (*) 1 9 Aleurone Color (Munsell code Lighter than | 1 2.5 Y 9/2) | | 1 9 (Munsell code Lighter Than 2.5 Y 9/2) | | |
| * 0 7 Hard Endosperm Color (Munsell code 2.5 Y | 8/10) | | 0 7 (Mui | nsell code 2.5 Y 8/10 |) |
| * 0 3 Endosperm Type: 1=Sweet (sul) 2=Extra Swe 4=High Amylose Starch 5=Waxy Starch 6=Hi 8=Super Sweet (se) 9=High Oil 10=Other | et (sh2) 3=Normal St gh Protein 7=High Ly | arch sine | 0 3 | | |
| 2 6. 6 gm Weight per 100 Kernels (unsized sample | 5.6 | 1700 seeds | 2 9 .7 | 8.7 | 1200 seeds |
| 9. COB: | Standard Deviation | Sample Size | | Standard Deviation | Sample Size |
| * 2 0.9 mm Cob Diameter at mid-point | 1.3 | 15 | 2 2.1 | 0.8 | 15 |
| 1 4 Cob Color (Munsell code 5 R 3/8) | | | 1 4 (Mi | unsell code 5 R 3/8) | |
| 10. DISEASE RESISTANCE (Rate from 1 (most susceptible) leave blank if not tested; leave Race or Strai | to 9 (most resistan n Options blank if p | :); olygenic): | | | |
| A. Leaf Blights, Wilts, and Local Infection Diseases | | | | | |
| 6 Anthracnose Leaf Blight (Colletotrichum graminicol. 7 Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) 7 Eyespot (Kabatiella zeae) 8 Goss's Wilt (Clavibacter michiganense spp. nebrask. 5 Gray Leaf Spot (Cercospora zeae-maydis) 7 Helminthosporium Leaf Spot (Bipolaris zeicola) Race 7 Northern Leaf Blight (Exserohilum turcicum) Race 1 8 Southern Leaf Blight (Bipolaris maydis) Race 0 Southern Rust (Puccinia polysora) 7 Stewart's Wilt (Erwinia stewartii) Other (Specify) | ense) e 2 | | 7 - - - - - - - - - - - - - | | |
| B. Systemic Diseases | | | | | |
| 6 Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Strain Sorghum Downy Mildew of Corn (Peronosclerospora soil Other (Specify) C. Stalk Rots | rghi) | | 3 - - - Strain - | · | |
| Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) | | | _ _ _ | | |
| D. Ear and Kernel Rots | | | | | |
| Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) |) | | | | |
| Application Variety Data | | | Standard I | nbred Data | |
| Note: Use chart on first page to choose color codes for | r color traits. | I | | | |

| | | | | 2005 00 | 111 |
|---|---------------------------|----------------------------------|--|-----------------------|----------------|
| Application Variety Data | Page 4 | | Standard Inbred Data | | |
| 11. INSECT RESISTANCE (Rate from 1 (most susceptible) to 9 (most blank if not tested): | ost resistar | nt); | | | |
| | andard viation | Sample Size | - | Standard Deviation | Sample Size |
| Silk Feeding: | | | _ | | |
| _ Ear Damage Corn Leaf Aphid (<i>Rhopalosiphum maidis</i>) | | | | ****************** | |
| Corn Sap Beetle (Carpophilus dimidiatus) European Corn Borer (Ostrinia nubilalis) 1 lst Generation (Typically Whorl Leaf Feeding) 2nd Generation (Typically Leaf Sheath-Collar Feeding) | | | _ _ 3 | | |
| Stalk Tunneling : . cm tunneled/plant | | | - | | |
| Fall Armyworm (Spodoptera frugiperda) Leaf Feeding | | | | | |
| Silk-Feeding : mg larval wt Maize Weevil (Sitophilus zeamaize) | | | | | |
| Northern Rootworm (Diabrotica barberi) Southern Rootworm (Diabrotica undecimpunctata) Southwestern Corn Borer (Diatraea grandiosella) Leaf Feeding | | | | | |
| Stalk Tunneling: cm tunneled/plant Two-spotted Spider Mite (Tetranychus urticae) Western Rootworm (Diabrotica virgifera virgifera) | <u>.</u> | | · _ | | |
| Other (Specify) 2. AGRONOMIC TRAITS: | | | | | |
| 7 Stay Green (at 65 days after anthesis) (Rate on a | scale from | 1=worst | 2 | | |
| to 9=excellent.) 0 0.0 % Dropped Ears (at 65 days after anthesis) | Douge III | 1 40100 | 0 0.0 | | |
| 0 0.0 % Pre-anthesis Brittle Snapping | | | 0 0.0 | | |
| 0 0.0 % Pre-anthesis Root Lodging | | | 0 0.0 | | |
| 0 0.0% Post-anthesis Root Lodging (at 65 days after an | thesis) | | 0 0.0 | | |
| Kg/ha Yield of Inbred Per Se (at 12-13% grain mo | isture) | | | | |
| 3. MOLECULAR MARKERS: (0=data unavailable; 1=data available b | ut not supp | lied; 2=data su | pplied) | | |
| 1 Isozymes 0 RFLP's 0 RAPD's | | | | | |
| EFERENCES: | | | | | |
| Butler, D.R. 1954. A System for the Classification of Corn In Emerson, R.A., G.W. Beadle, and A.C. Fraser. 1935. A Summary Farr, D.F., G.F. Bills, G.P. Chamuris, A.Y. Rossman. 1989. Fu Phytopathological Society, St. Paul, MN. Inglett, G.E. (Ed.) 1970. Corn: Culture, Processing, Products | of Linkage ngi on Plan | Studies in Maiz and Plant Pro | e. Cornell A.E.S. ducts in the Unit | . Mem. 180. | merican |

Inglett, G.E. (Ed.) 1970. Corn: Culture, Processing, Products. Avi Publishing Company, Westport, CT.
Jugenheimer, R.W. 1976. Corn: Improvement, Seed Production, and Uses. John Wiley & Sons, New York.
McGee, D.C. 1988. Maize Diseases. APS Press, St. Paul, MN. 150 pp.
Munsell Color Chart for Plant Tissues. Macbeth. P.O. Box 230. Newburgh, N.Y. 12551-0230
The Mutants of Maize. 1968. Crop Science Society of America. Madison, WI.
Shurtleff, M.C. 1980. Compendium of Corn Diseases. APS Press, St. Paul, MN. 105 pp.
Sprague, G.F., and J.W. Dudley (Editors). 1988. Corn and Corn Improvement, Third Edition. Agronomy Monograph 18. ASA, CSSA, Madison, WI.

Madison, WI.
Stringfield, G.H. Maize Inbred Lines of Ohio. Ohio A.E.S., Bul. 831. 1959.
U.S. Department of Agriculture. 1936, 1937. Yearbook.

COMMENTS (eg. state how heat units were calculated, standard inbred seed source, and/or where data was collected. Continue in Exhibit

Heat Unit Calculation: GDU = Daily Max Temp ($<=86^{\circ}F$) + Daily Min Temp ($>=50^{\circ}F$) - $50^{\circ}F$

Supplemental data provided for pollen shed, ear weight, % round kernels and weight per 100 kernels from 2006 production parent test data and 2006 seed inventory data.

| REPRODUCE LOCALLY. Include form number and edition date on a | all reproductions. | FORM APPROVED - OMB No. 0581-0055 |
|--|---|---|
| U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E | Application is required in order to de certificate is to be issued (7 U.S.C. 2 | etermine if a plant variety protection |
| STATEMENT OF THE BASIS OF OWNERSHIP | confidential until the certificate is iss | ued (7 U.S.C. 2426) |
| 1. NAME OF APPLICANT(S) | 2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER | 3. VARIETY NAME |
| Monsanto Technology L.L.C. | | 1006234 |
| 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country) | 5. TELEPHONE (Include area code) | 6. FAX (Include area code) |
| 800 N. Lindbergh Blvd. | (815) 758-9281 | (815) 758-3117 |
| Creve Couer, MO 63167 | 7. PVPO NUMBER | |
| U.S.A. | | 200500157 |
| 8. Does the applicant own all rights to the variety? Mark an "X" in the | e appropriate block. If no, please expla | in. X YES NO |
| | | |
| | | |
| | • | : |
| 9. Is the applicant (individual or company) a U.S. national or a U.S. b | ased company? If no, give name of co | ountry. YES NO |
| | | |
| 10. Is the applicant the original owner? | NO If no, please answer one | of the following: |
| | · | |
| a. If the original rights to variety were owned by individual(s), is (| | |
| L 1E3 | NO If no, give name of country | у |
| | | |
| b. If the original rights to variety were owned by a company(ies), | is (are) the original owner(s) a U.S. base | ed company? |
| YES | NO If no, give name of country | y . |
| | | |
| 11. Additional explanation on ownership (Trace ownership from origin | al breeder to current owner. Use the re | verse for extra space if needed): |
| | | |
| Corn Variety I006234 was originated and dev | reloned by a breeder employed | hy Managata |
| Technology L.L.C. By agreement between M | lonsanto Technology L.L.C. and | the breeder, all |
| rights to any invention, discovery or develope | nent are assigned to Monsanto | Technology I. J. C. |
| No rights to such invention, discovery or deve | elopment are retained by the bre | eder. |
| | | |
| PLEASE NOTE: | | |
| | | |
| Plant variety protection can only be afforded to the owners (not license | es) who meet the following criteria: | |
| If the rights to the variety are owned by the original breeder, that per national of a country which affords similar protection to nationals of t | son must be a U.S. national, national of the U.S. for the same genus and species | a UPOV member country, or s. |
| If the rights to the variety are owned by the company which employe nationals of a UPOV member country, or owned by nationals of a co genus and species. | nd the original breeder(s), the company rountry which affords similar protection to | nust be U.S. based, owned by nationals of the U.S. for the same |
| . If the applicant is an owner who is not the original owner, both the or | iginal owner and the applicant must me | et one of the above criteria. |
| the original breeder/owner may be the individual or company who directly the company which is the company when the company which is the company which is the company which is the company when the company which is the co | | |
| | | |

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.